

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 October 2003 (16.10.2003)

PCT

(10) International Publication Number
WO 03/085137 A1

(51) International Patent Classification⁷: C12Q 1/68,
A01N 43/04, C07H 21/04, A61K 31/07

(21) International Application Number: PCT/US03/09982

(22) International Filing Date: 1 April 2003 (01.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/114,544 1 April 2002 (01.04.2002) US

(71) Applicant (*for all designated States except US*): ISIS
PHARMACEUTICALS, INC [US/US]; 2292 Faraday
Avenue, Carlsbad, CA 92008 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): MONIA, Brett, P.
[US/US]; 2306 Casa Hermosa Court, Encinitas, CA 92024
(US). COWSERT, Lex, M. [US/US]; 2367 West Gate
Drive, Pittsburgh, PA 15237 (US).

(74) Agent: WILLIAMS, Joseph, A., Jr.; Marshall Gerstein &
Borun, 233 S. Wacker Drive, 6300 Sears Tower, Chicago,
IL 60606-6357 (US).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

ANTISENSE MODULATION OF LIVER GLYCOGEN PHOSPHORYLASE EXPRESSION

(57) Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of liver glycogen phosphorylase. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding liver glycogen phosphorylase. Methods of using these compounds for modulation of liver glycogen phosphorylase expression.

5 **ANTISENSE MODULATION OF LIVER**
 GLYCOGEN PHOSPHORYLASE EXPRESSION

10 **FIELD OF THE INVENTION**

15 The present invention provides compositions and methods for modulating the expression of liver glycogen phosphorylase. In particular, this invention relates to antisense compounds, particularly oligonucleotides, specifically hybridizable with nucleic acids encoding human liver glycogen phosphorylase. Such oligonucleotides have been shown to modulate the expression of liver glycogen phosphorylase.

20 **BACKGROUND OF THE INVENTION**

25 There are several mechanisms in place for the homeostatic regulation of biological processes, and balanced energy metabolism is critical to all of these mechanisms. In higher organisms energy stores are in the form of glycogen and upon energy deficit these stores are mobilized through enzymatic digestion to glucose-1-phosphate by the glycogen phosphorylase family of proteins.

30 Mammalian glycogen phosphorylases comprise a family of three isozymes which are distinguished by their

-2-

electrophoretic mobilities, immunological properties and tissue-specific distribution. Each isozyme is encoded by a different gene, and these genes have been denoted PYGL, PYGM and PYGB, for liver, muscle and brain isoforms, respectively (Newgard et al., *Crit. Rev. Biochem. Mol. Biol.*, 1989, 24, 69-99). The primary control however, common to all isozymes, is the phosphorylation of the inactive state, (b), to the active phosphorylated state, (a). This phosphorylation on serine-14 stabilizes the subunits of the homodimer and alters the binding sites for allosteric effectors and substrates (Sprang et al., *Nature*, 1988, 336, 215-221).

Liver glycogen phosphorylase (also known as 1,4- α -D-glucan:orthophosphate α -D-glucosyltransferase, glycogen phosphorylase (liver), EC 2.4.1.1 and HLGPa, for human liver glycogen phosphorylase a) is the enzyme which catalyzes the degradation of stored glycogen in the liver to glucose-1-phosphate by the cleavage of α -1,4-glycosidic bonds and therefore plays a critical role in carbohydrate metabolism and blood glucose homeostasis (Newgard et al., *Proc. Natl. Acad. Sci. U. S. A.*, 1986, 83, 8132-8136). The activity of liver glycogen phosphorylase is tightly regulated requiring the presence of a cofactor, pyridoxal phosphate, and involving allosteric mechanisms which include activation by AMP and glycogen binding and inhibition by glucose and glucose-6-phosphate binding. The enzyme is also regulated through phosphorylation by phosphorylase kinase which activates the homodimer (Keppens et al., *Hepatology*, 1993, 17, 610-614).

The gene for liver glycogen phosphorylase (PYGL) has been mapped to chromosome 14 and mutations in this gene give rise to glycogen storage disease type VI (GSD VI) or Hers Disease, a group of disorders that cause hepatomegaly.

and hypoglycemia (Burwinkel et al., *Am. J. Hum. Genet.*, 1998, 62, 785-791; Chang et al., *Hum. Mol. Genet.*, 1998, 7, 865-870; Newgard et al., *Am. J. Hum. Genet.*, 1987, 40, 351-364). These mutations consist of two splice-site mutations which result in aberrant exon retention and exon skipping and two missense mutations which produce nonconservative replacements of amino acids that are normally conserved in all eukaryotes.

To date, two types of inhibitors targeting glycogen phosphorylase function have been reported. These involved the use of glucose analogs containing multiple polar groups which bind near the active site of the protein (Lundgren et al., 1997; Lundgren and Kirk, 1995) and caffeine and other heteroaromatic analogs which bind at the purine inhibitory site (Kasvinsky et al., *Can. J. Biochem.*, 1981, 59, 387-395; Kasvinsky et al., *J. Biol. Chem.*, 1978, 253, 3343-3351; Kasvinsky et al., *J. Biol. Chem.*, 1978, 253, 9102-9106). However, none of these compounds have been shown to be orally active, limiting their utility.

Recently, the discovery of an orally active compound, CP-91149 and derivatives thereof, that lowers plasma glucose levels in an animal model of type 2 diabetes was reported (Hoover et al., *J. Med. Chem.*, 1998, 41, 2934-2938; Martin et al., *Proc. Natl. Acad. Sci. U. S. A.*, 1998, 95, 1776-1781). This indole-containing compound was shown to inhibit glycogenolysis in diabetic ob/ob mice, and in rat and human liver cells by inhibiting liver glycogen phosphorylase. It is believed that inhibition of glycogenolysis will be of therapeutic benefit in the treatment of diabetes, particularly type II diabetes.

Antisense technology is emerging as an effective means for reducing the expression of specific gene products and may therefore prove to be uniquely useful in a number of

-4-

therapeutic, diagnostic, and research applications for the modulation of glycogen phosphorylase expression.

SUMMARY OF THE INVENTION

5 The present invention is directed to antisense compounds, particularly oligonucleotides, which are targeted to a nucleic acid encoding liver glycogen phosphorylase, and which modulate the expression of liver glycogen phosphorylase. Pharmaceutical and other
10 compositions comprising the antisense compounds of the invention are also provided. Further provided are methods of modulating the expression of liver glycogen phosphorylase in cells or tissues comprising contacting said cells or tissues with one or more of the antisense
15 compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of liver glycogen phosphorylase by administering a therapeutically or
20 prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

DETAILED DESCRIPTION OF THE INVENTION

 The present invention employs oligomeric antisense
25 compounds, particularly oligonucleotides, for use in modulating the function of nucleic acid molecules encoding liver glycogen phosphorylase, ultimately modulating the amount of liver glycogen phosphorylase produced. This is accomplished by providing antisense compounds which
30 specifically hybridize with one or more nucleic acids encoding liver glycogen phosphorylase. As used herein, the terms "target nucleic acid" and "nucleic acid encoding liver glycogen phosphorylase" encompass DNA encoding liver glycogen phosphorylase, RNA (including pre-mRNA and mRNA)

transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of liver glycogen phosphorylase. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding liver glycogen phosphorylase. The targeting process also includes determination of a site or sites within this gene for the

-6-

antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA molecule transcribed from a gene encoding liver glycogen phosphorylase, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of

-7-

such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to

- 8 -

form a continuous mRNA sequence. mRNA splice sites, i.e., intron-exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions are also preferred targets. It has also been found that introns can also be effective, and therefore preferred, target regions for antisense compounds targeted, for example, to DNA or pre-mRNA.

Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds.

"Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing.

such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

Antisense compounds are commonly used as research reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotides have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment

-10-

regimes for treatment of cells, tissues and animals, especially humans.

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 30 nucleobases (i.e. from about 8 to about 30 linked nucleosides). Particularly preferred antisense compounds are antisense oligonucleotides, even more preferably those comprising from about 12 to about 25 nucleobases. As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In

-11-

forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkyl-phosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are

-12-

also included.

Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808;
5 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897;
5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131;
5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677;
5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111;
5,563,253; 5,571,799; 5,587,361; and 5,625,050, certain of
10 which are commonly owned with this application, and each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl
15 internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside);
20 siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide
25 backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444;
30 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564;
5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677;
5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289;
5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070;
5,663,312; 5,633,360; 5,677,437; and 5,677,439, certain of

-13-

which are commonly owned with this application, and each of which is herein incorporated by reference.

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to azanitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497-1500.

Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular $-\text{CH}_2-\text{NH}-\text{O}-\text{CH}_2-$, $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{O}-\text{CH}_2-$ [known as a methylene (methylimino) or MMI backbone], $-\text{CH}_2-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-$, $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{N}(\text{CH}_3)-\text{CH}_2-$ and $-\text{O}-\text{N}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-$ [wherein the native phosphodiester backbone is represented as $-\text{O}-\text{P}-\text{O}-\text{CH}_2-$] of the above referenced U.S. patent 5,489,677, and the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. patent 5,034,506.

Modified oligonucleotides may also contain one or more

-14-

substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Particularly preferred are O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. Other preferred oligonucleotides

comprise one of the following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., *Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoethoxy, i.e., a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH₂-O-CH₂-N(CH₂)₂, also described in examples hereinbelow.

Other preferred modifications include 2'-methoxy (2'-O-CH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in

-15-

2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents
5 that teach the preparation of such modified sugar structures include, but are not limited to, U.S.:
4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878;
5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811;
5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053;
10 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

Oligonucleotides may also include nucleobase (often
15 referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other
20 synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-
25 thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl
30 and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in United States Patent No.

-16-

3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; and 5,681,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference, and United States patent 5,750,692, which is commonly owned with the instant application and also herein incorporated by reference.

-17-

Another modification of the oligonucleotides of the invention involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660, 306-309; Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10, 1111-1118; Kabanov et al., *FEBS Lett.*, 1990, 259, 327-330; Svinarchuk et al., *Biochimie*, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654; Shea et al., *Nucl. Acids Res.*, 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277, 923-937).

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717, 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802;

-18-

5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718;
5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779;
4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582;
4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830;
5 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506;
5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241,
5,391,723; 5,416,203, 5,451,463; 5,510,475; 5,512,667;
5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481;
5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and
10 5,688,941, certain of which are commonly owned with the
instant application, and each of which is herein
incorporated by reference.

It is not necessary for all positions in a given
compound to be uniformly modified, and in fact more than
15 one of the aforementioned modifications may be incorporated
in a single compound or even at a single nucleoside within
an oligonucleotide. The present invention also includes
antisense compounds which are chimeric compounds.
"Chimeric" antisense compounds or "chimeras," in the
20 context of this invention, are antisense compounds,
particularly oligonucleotides, which contain two or more
chemically distinct regions, each made up of at least one
monomer unit, i.e., a nucleotide in the case of an
oligonucleotide compound. These oligonucleotides typically
25 contain at least one region wherein the oligonucleotide is
modified so as to confer upon the oligonucleotide increased
resistance to nuclease degradation, increased cellular
uptake, and/or increased binding affinity for the target
nucleic acid. An additional region of the oligonucleotide
30 may serve as a substrate for enzymes capable of cleaving
RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is
a cellular endonuclease which cleaves the RNA strand of an
RNA:DNA duplex. Activation of RNase H, therefore, results
in cleavage of the RNA target, thereby greatly enhancing

-19-

the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

Chimeric antisense compounds of the invention may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the phosphorothioates and alkylated derivatives.

The antisense compounds of the invention are synthesized in vitro and do not include antisense compositions of biological origin, or genetic vector

-20-

constructs designed to direct the in vivo synthesis of antisense molecules.

The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption assisting formulations include, but are not limited to, U.S.:
5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291;
5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330;
4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170;
15 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978;
5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259;
5,543,152; 5,556,948; 5,580,575; and 5,595,756, each of which is herein incorporated by reference.

The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as ~~SATS~~.

[(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510 to Gosselin et al., published December 9, 1993 or in WO 94/26764 to Imbach et al.

5 The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

10 Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are

15 N,N'-dibenzylethylenediamine, chloroprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge et al., "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-19). The base addition salts of said acidic compounds
20 are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by contacting the salt form with an acid and isolating the free acid in the conventional manner. The
25 free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for purposes of the present invention. As used herein, a "pharmaceutical addition
30 salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and

-22-

phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

For oligonucleotides, preferred examples of pharmaceutically acceptable salts include but are not

-23-

limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from elemental anions such as chlorine, bromine, and iodine.

The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of liver glycogen phosphorylase is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding liver glycogen phosphorylase, enabling sandwich and other assays to easily be constructed to exploit this fact. Hybridization of the

-24-

antisense oligonucleotides of the invention with a nucleic acid encoding liver glycogen phosphorylase can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of liver glycogen phosphorylase in a sample may also be prepared.

The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

Compositions and formulations for oral administration

-25-

include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets or tablets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable.

5 Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and
10 other pharmaceutically acceptable carriers or excipients.

Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions may be generated from a variety of components that include, but
15 are not limited to, preformed liquids, self-emulsifying solids and self-emulsifying semisolids.

The pharmaceutical formulations of the present invention, which may conveniently be presented in unit dosage form, may be prepared according to conventional
20 techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general the formulations are prepared by uniformly and intimately bringing into association the
25 active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

The compositions of the present invention may be formulated into any of many possible dosage forms such as,
30 but not limited to, tablets, capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity

-26-

of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

15

Emulsions

The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1 μm in diameter. (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 301). Emulsions are often biphasic systems comprising of two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be either water-in-oil (w/o) or of the oil-in-water (o/w) variety. When an aqueous phase is

-27-

finely divided into and dispersed as minute droplets into a bulk oily phase the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase the resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water stabilized in an oily continuous provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories:

synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal hydroxides, nonswelling clays such as

-29-

bentonite, attapulgite, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

5 A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants
10 (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

15 Hydrophilic colloids or hydrocolloids include naturally occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose
20 and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase
25 droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these
30 formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added

-30-

to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of reasons of ease of formulation, efficacy from an absorption and bioavailability standpoint. (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and

-31-

then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 271).

The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers,

-32-

polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art.

The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides

(Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug

absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides et al., *Pharmaceutical Research*, 1994, 11, 1385; Ho et al., *J. Pharm. Sci.*, 1996, 85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

-34-

Liposomes

There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which is highly deformable and able to pass through such fine pores.

Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important considerations in the preparation of liposome

-35-

formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

Liposomes are useful for the transfer and delivery of active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes. As the merging of the liposome and cell progresses, the liposomal contents are emptied into the cell where the active agent may act.

Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the skin. The majority of applications resulted in the targeting of the upper epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into

-36-

the cell cytoplasm (Wang et al., *Biochem. Biophys. Res. Commun.*, 1987, 147, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou et al., *Journal of Controlled Release*, 1992, 19, 269-274).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (e.g. as a solution or as an emulsion) were ineffective (Weiner et al., *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal

-37-

formulation was superior to aqueous administration (du Plessis et al., *Antiviral Research*, 1992, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu et al. *S.T.P. Pharma. Sci.*, 1994, 4, 6, 466).

Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G_{M1}, or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen et al., *FEBS Letters*, 1987, 223, 42; Wu et al.,

-38-

Cancer Research, 1993, 53, 3765). Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos et al. (Ann. N.Y. Acad. Sci., 1987, 507, 64) reported the ability of monosialoganglioside G_{M1} ,
5 galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. These findings were expounded upon by Gabizon et al. (Proc. Natl. Acad. Sci. U.S.A., 1988, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen et al., disclose liposomes
10 comprising (1) sphingomyelin and (2) the ganglioside G_{M1} or a galactocerebroside sulfate ester. U.S. Patent No. 5,543,152 (Webb et al.) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-sn-dimyristoylphosphatidylcholine are disclosed in WO 97/13499
15 (Lim et al.).

Many liposomes comprising lipids derivatized with one or more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto et al. (Bull. Chem. Soc. Jpn., 1980, 53, 2778) described liposomes
20 comprising a nonionic detergent, 2C₁₂15G, that contains a PEG moiety. Illum et al. (FEBS Lett., 1984, 167, 79) noted that hydrophilic coating of polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the
25 attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Patent Nos. 4,426,330 and 4,534,899). Klivanov et al. (FEBS Lett., 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE)
30 derivatized with PEG or PEG stearate have significant increases in blood circulation half-lives. Blume et al. (Biochimica et Biophysica Acta, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids.

-39-

e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle et al. (U.S. Patent Nos. 5,013,556 and 5,356,633) and Martin et al. (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin et al.) and in WO 94/20073 (Zalipsky et al.). Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi et al.). U.S. Patent Nos. 5,540,935 (Miyazaki et al.) and 5,556,948 (Tagawa et al.) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry et al. discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to Tagawa et al. discloses protein-bonded liposomes and asserts that the contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman et al. describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love et al. discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable that they are easily able to penetrate through

-40-

pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in this class. The

-41-

polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

If the surfactant molecule has the ability to carry either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations and in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized and nonionized forms. However, usually

-42-

only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid; with the result that absorption of oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi et al., *J. Pharm. Pharmacol.*, 1988, 40, 252).

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid,

-43-

- linolenic acid, dicaprinate, tricaprinate, monoolein (1-monooleoyl-rac-glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprinate, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C₁₋₁₀ alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and di-glycerides thereof (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-654).

- Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucolic acid (sodium glucolate), glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-24,25-dihydrofusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical*

-44-

- Reviews in Therapeutic Drug Carrier Systems, 1991, page 92; Swinyard, Chapter 39 In: Remington's Pharmaceutical Sciences, 18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, Critical
- 5 Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1-33; Yamamoto et al., J. Pharm. Exp. Ther., 1992, 263, 25; Yamashita et al., J. Pharm. Sci., 1990, 79, 579-583).

- Chelating Agents: Chelating agents, as used in
- 10 connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration
- 15 enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, J. Chromatogr., 1993, 618, 315-339).
- 20 Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), N-acyl derivatives of collagen, laureth-9 and N-amino acyl derivatives of beta-
- 25 diketones (enamines) (Lee et al., Critical Reviews in Therapeutic Drug Carrier Systems, 1991, page 92; Muranishi, Critical Reviews in Therapeutic Drug Carrier Systems, 1990, 7, 1-33; Buur et al., J. Control Rel., 1990, 14, 43-51).

- 30 Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as ~~chelating agents or as surfactants but that~~

nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33). This class of penetration enhancers include, for example, 5 unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., 10 *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. 15 Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the 20 penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

25 Carriers

Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (i.e., does 30 not possess biological activity per se) but is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation.

-46-

The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'isothiocyano-stilbene-2,2'-disulfonic acid (Miyao et al., *Antisense Res. Dev.*, 1995, 5, 115-121; Takakura et al., *Antisense & Nucl. Acid Drug Dev.*, 1996, 6, 177-183).

15 Excipients

In contrast to a carrier compound, a "pharmaceutical carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (e.g., pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose, etc.); fillers (e.g., lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, etc.); lubricants (e.g., magnesium stearate, talc, silica, colloidal silicon dioxide, stearic acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium

acetate, etc.); disintegrants (e.g., starch, sodium starch glycolate, etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic
5 excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions,
10 alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous
15 solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for
20 non-parenteral administration which do not deleteriously react with nucleic acids can be used.

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose,
25 magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Other Components

The compositions of the present invention may
30 additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example,

-48-

antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as
5 dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations
10 can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact
15 with the nucleic acid(s) of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

20 Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents
25 include, but are not limited to, anticancer drugs such as daunorubicin, dactinomycin, doxorubicin, bleomycin, mitomycin, nitrogen mustard, chlorambucil, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine (CA), 5-fluorouracil (5-FU), floxuridine
30 (5-FUdR), methotrexate (MTX), colchicine, vincristine, vinblastine, etoposide, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 1206-1228). Anti-inflammatory

-49-

drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in
5 compositions of the invention. See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more
10 combined compounds may be used together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds
15 targeted to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their subsequent administration is believed to be within the
20 skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules
25 can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and
30 can generally be estimated based on EC_{50} s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years.

-50-

Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

EXAMPLES

Example 1

Nucleoside Phosphoramidites for Oligonucleotide Synthesis

Deoxy and 2'-alkoxy amidites

2'-Deoxy and 2'-methoxy beta-cyanoethyldiisopropyl phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA). Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, the standard cycle for unmodified oligonucleotides was utilized, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds.

Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-C) nucleotides were synthesized according to published methods [Sanghvi, et. al., *Nucleic Acids Research*, 1993, 21, 3197-3203] using commercially available

-51-

phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA).

2'-Fluoro amidites

5 2'-Fluorodeoxyadenosine amidites

2'-fluoro oligonucleotides were synthesized as described previously [Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841] and United States patent 5,670,633, herein incorporated by reference. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and by modifying literature procedures whereby the 2'-alpha-fluoro atom is introduced by a S_N2-displacement of a 2'-beta-trityl group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies and standard methods were used to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

2'-Fluorodeoxyguanosine

The synthesis of 2'-deoxy-2'-fluoroguanosine was accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate diisobutryl-arabinofuranosylguanosine. Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP to give diisobutryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies were used to obtain the 5'-DMT- and

-52-

5'-DMT-3'-phosphoramidites.

2'-Fluorouridine

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

10

2'-Fluorodeoxycytidine

2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

15

2'-O-(2-Methoxyethyl) modified amidites

2'-O-Methoxyethyl-substituted nucleoside amidites are prepared as follows, or alternatively, as per the methods of Martin, P., *Helvetica Chimica Acta*, 1995, 78, 486-504.

20

2,2'-Anhydro[1-(beta-D-arabinofuranosyl)-5-methyluridine]

25 5-Methyluridine (ribosylthymine, commercially available through Yamasa, Choshi, Japan) (72.0 g, 0.279 M), diphenylcarbonate (90.0 g, 0.420 M) and sodium bicarbonate (2.0 g, 0.024 M) were added to DMF (300 mL). The mixture was heated to reflux, with stirring, allowing the evolved carbon dioxide gas to be released in a controlled manner. After 1 hour, the slightly darkened solution was concentrated under reduced pressure. The resulting syrup was poured into diethylether (2.5 L), with stirring. The product formed a gum. The ether was decanted and the

30

-53-

residue was dissolved in a minimum amount of methanol (ca. 400 mL). The solution was poured into fresh ether (2.5 L) to yield a stiff gum. The ether was decanted and the gum was dried in a vacuum oven (60°C at 1 mm Hg for 24 h) to give a solid that was crushed to a light tan powder (57 g, 85% crude yield). The NMR spectrum was consistent with the structure, contaminated with phenol as its sodium salt (ca. 5%). The material was used as is for further reactions (or it can be purified further by column chromatography using a gradient of methanol in ethyl acetate (10-25%) to give a white solid, mp 222-4°C).

2'-O-Methoxyethyl-5-methyluridine

2,2'-Anhydro-5-methyluridine (195 g, 0.81 M), tris(2-methoxyethyl)borate (231 g, 0.98 M) and 2-methoxyethanol (1.2 L) were added to a 2 L stainless steel pressure vessel and placed in a pre-heated oil bath at 160°C. After heating for 48 hours at 155-160°C, the vessel was opened and the solution evaporated to dryness and triturated with MeOH (200 mL). The residue was suspended in hot acetone (1 L). The insoluble salts were filtered, washed with acetone (150 mL) and the filtrate evaporated. The residue (280 g) was dissolved in CH₃CN (600 mL) and evaporated. A silica gel column (3 kg) was packed in CH₂Cl₂/acetone/MeOH (20:5:3) containing 0.5% Et₃NH. The residue was dissolved in CH₂Cl₂ (250 mL) and adsorbed onto silica (150 g) prior to loading onto the column. The product was eluted with the packing solvent to give 160 g (63%) of product. Additional material was obtained by reworking impure fractions.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine

2'-O-Methoxyethyl-5-methyluridine (160 g, 0.506 M) was co-evaporated with pyridine (250 mL) and the dried residue

-54-

dissolved in pyridine (1.3 L). A first aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the mixture stirred at room temperature for one hour. A second aliquot of dimethoxytrityl chloride (94.3 g, 0.278 M) was added and the reaction stirred for an additional one hour. Methanol (170 mL) was then added to stop the reaction. HPLC showed the presence of approximately 70% product. The solvent was evaporated and triturated with CH₃CN (200 mL). The residue was dissolved in CHCl₃ (1.5 L) and extracted with 2x500 mL of saturated NaHCO₃ and 2x500 mL of saturated NaCl. The organic phase was dried over Na₂SO₄, filtered and evaporated. 275 g of residue was obtained. The residue was purified on a 3.5 kg silica gel column, packed and eluted with EtOAc/hexane/acetone (5:5:1) containing 0.5% Et₃NH. The pure fractions were evaporated to give 164 g of product. Approximately 20 g additional was obtained from the impure fractions to give a total yield of 183 g (57%).

20 **3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (106 g, 0.167 M), DMF/pyridine (750 mL of a 3:1 mixture prepared from 562 mL of DMF and 188 mL of pyridine) and acetic anhydride (24.38 mL, 0.258 M) were combined and stirred at room temperature for 24 hours. The reaction was monitored by TLC by first quenching the TLC sample with the addition of MeOH. Upon completion of the reaction, as judged by TLC, MeOH (50 mL) was added and the mixture evaporated at 35°C. The residue was dissolved in CHCl₃ (800 mL) and extracted with 2x200 mL of saturated sodium bicarbonate and 2x200 mL of saturated NaCl. The water layers were back extracted with 200 mL of CHCl₃. The combined organics were dried with sodium sulfate and evaporated to give 122 g of residue (approx. 90% product).

-55-

The residue was purified on a 3.5 kg silica gel column and eluted using EtOAc/hexane(4:1). Pure product fractions were evaporated to yield 96 g (84%). An additional 1.5 g was recovered from later fractions.

5

3'-O-Acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine

A first solution was prepared by dissolving 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyluridine (96 g, 0.144 M) in CH₃CN (700 mL) and set
10 aside. Triethylamine (189 mL, 1.44 M) was added to a solution of triazole (90 g, 1.3 M) in CH₃CN (1 L), cooled to -5°C and stirred for 0.5 h using an overhead stirrer. POCl₃ was added dropwise, over a 30 minute period, to the stirred
15 solution maintained at 0-10°C, and the resulting mixture stirred for an additional 2 hours. The first solution was added dropwise, over a 45 minute period, to the latter solution. The resulting reaction mixture was stored overnight in a cold room. Salts were filtered from the
20 reaction mixture and the solution was evaporated. The residue was dissolved in EtOAc (1 L) and the insoluble solids were removed by filtration. The filtrate was washed with 1x300 mL of NaHCO₃ and 2x300 mL of saturated NaCl, dried over sodium sulfate and evaporated. The residue was
25 triturated with EtOAc to give the title compound.

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine

A solution of 3'-O-acetyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methyl-4-triazoleuridine (103 g, 0.141 M)
30 in dioxane (500 mL) and NH₄OH (30 mL) was stirred at room temperature for 2 hours. The dioxane solution was ~~evaporated and the residue azeotroped with MeOH (2x200 mL).~~ The residue was dissolved in MeOH (300 mL) and transferred

-56-

to a 2 liter stainless steel pressure vessel. MeOH (400 mL) saturated with NH₃ gas was added and the vessel heated to 100°C for 2 hours (TLC showed complete conversion). The vessel contents were evaporated to dryness and the residue was dissolved in EtOAc (500 mL) and washed once with saturated NaCl (200 mL). The organics were dried over sodium sulfate and the solvent was evaporated to give 85 g (95%) of the title compound.

10 **N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine**

2'-O-Methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (85 g, 0.134 M) was dissolved in DMF (800 mL) and benzoic anhydride (37.2 g, 0.165 M) was added with stirring. After stirring for 3 hours, TLC showed the reaction to be approximately 95% complete. The solvent was evaporated and the residue azeotroped with MeOH (200 mL). The residue was dissolved in CHCl₃ (700 mL) and extracted with saturated NaHCO₃ (2x300 mL) and saturated NaCl (2x300 mL), dried over MgSO₄ and evaporated to give a residue (96 g). The residue was chromatographed on a 1.5 kg silica column using EtOAc/hexane (1:1) containing 0.5% Et₃NH as the eluting solvent. The pure product fractions were evaporated to give 90 g (90%) of the title compound.

25

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine-3'-amidite

N4-Benzoyl-2'-O-methoxyethyl-5'-O-dimethoxytrityl-5-methylcytidine (74 g, 0.10 M) was dissolved in CH₂Cl₂ (1 L). Tetrazole diisopropylamine (7.1 g) and 2-cyanoethoxy-tetra(isopropyl)phosphite (40.5 mL, 0.123 M) were added with stirring, under a nitrogen atmosphere. The resulting mixture was stirred for 20 hours at room temperature (TLC showed the reaction to be 95% complete). The reaction

-57-

mixture was extracted with saturated NaHCO_3 (1x300 mL) and saturated NaCl (3x300 mL). The aqueous washes were back-extracted with CH_2Cl_2 (300 mL), and the extracts were combined, dried over MgSO_4 and concentrated. The residue
5 obtained was chromatographed on a 1.5 kg silica column using EtOAc/hexane (3:1) as the eluting solvent. The pure fractions were combined to give 90.6 g (87%) of the title compound.

10 2'-O-(Aminooxyethyl) nucleoside amidites and 2'-O-(dimethylaminooxyethyl) nucleoside amidites

2'-(Dimethylaminooxyethoxy) nucleoside amidites

2'-(Dimethylaminooxyethoxy) nucleoside amidites [also
15 known in the art as 2'-O-(dimethylaminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected
20 with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

5'-O-tert-Butyldiphenylsilyl-O²-2'-anhydro-5-methyluridine

25 O²-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. tert-Butyldiphenylchlorosilane
30 (125.8g, 119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (R_f 0.22, ethyl acetate) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned

-58-

between dichloromethane (1 L) and saturated sodium bicarbonate (2x1 L) and brine (1 L). The organic layer was dried over sodium sulfate and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of ethyl acetate and ethyl ether (600mL) and the solution was cooled to -10°C. The resulting crystalline product was collected by filtration, washed with ethyl ether (3x200 mL) and dried (40°C, 1mm Hg, 24 h) to 149g (74.8%) of white solid. TLC and NMR were consistent with pure product.

5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine

In a 2 L stainless steel, unstirred pressure reactor was added borane in tetrahydrofuran (1.0 M, 2.0 eq, 622 mL). In the fume hood and with manual stirring, ethylene glycol (350 mL, excess) was added cautiously at first until the evolution of hydrogen gas subsided. 5'-O-tert-Butyldiphenylsilyl-O²-2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160 °C was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient and opened. TLC (R_f 0.67 for desired product and R_f 0.82 for ara-T side product, ethyl acetate) indicated about 70% conversion to the product. In order to avoid additional side product formation, the reaction was stopped, concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. [Alternatively, once the low boiling solvent is gone, the remaining solution can be partitioned between ethyl acetate and water. The product will be in the organic phase.] The residue was

-59-

purified by column chromatography (2kg silica gel, ethyl acetate-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, stripped and dried to product as a white crisp foam (84g, 50%), contaminated starting material (17.4g) and pure reusable starting material 20g. The yield based on starting material less pure recovered starting material was 58%. TLC and NMR were consistent with 99% pure product.

10 2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine

5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol). It was then dried over P₂O₅ under high vacuum for two days at 40°C. The reaction mixture was flushed with argon and dry THF (369.8mL, Aldrich, sure seal bottle) was added to get a clear solution. Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture. The rate of addition is maintained such that resulting deep red coloration is just discharged before adding the next drop. After the addition was complete, the reaction was stirred for 4 hrs. By that time TLC showed the completion of the reaction (ethylacetate:hexane, 60:40). The solvent was evaporated in vacuum. Residue obtained was placed on a flash column and eluted with ethyl acetate:hexane (60:40), to get 2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%).

-60-

5'-O-tert-butylldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine

2'-O-([2-phthalimidooxy)ethyl]-5'-t-butylldiphenylsilyl-5-methyluridine (3.1g, 4.5mmol) was dissolved in dry CH₂Cl₂ (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h the mixture was filtered, the filtrate was washed with ice cold CH₂Cl₂ and the combined organic phase was washed with water, brine and dried over anhydrous Na₂SO₄. The solution was concentrated to get 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). To this formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. Solvent was removed under vacuum; residue chromatographed to get 5'-O-tert-butylldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine as white foam (1.95 g, 78%).

5'-O-tert-Butylldiphenylsilyl-2'-O-[N,N-dimethylaminooxyethyl]-5-methyluridine

5'-O-tert-butylldiphenylsilyl-2'-O-[(2-formadoximinooxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium p-toluenesulfonate (PPTS) in dry MeOH (30.6mL). Sodium cyanoborohydride (0.39g, 6.13mmol) was added to this solution at 10°C under inert atmosphere. The reaction mixture was stirred for 10 minutes at 10°C. After that the reaction vessel was removed from the ice bath and stirred at room temperature for 2 h, the reaction monitored by TLC (5% MeOH in CH₂Cl₂). Aqueous NaHCO₃ solution (5%, 10mL) was added and extracted with ethyl acetate (2x20mL). Ethyl acetate phase was dried over anhydrous Na₂SO₄, evaporated to dryness. Residue was dissolved in a solution of 1M PPTS in MeOH (30.6mL). Formaldehyde (20% w/w, 30mL, 3.37mmol) was

-61-

added and the reaction mixture was stirred at room temperature for 10 minutes. Reaction mixture cooled to 10°C in an ice bath, sodium cyanoborohydride (0.39g, 6.13mmol) was added and reaction mixture stirred at 10°C for 10 minutes. After 10 minutes, the reaction mixture was removed from the ice bath and stirred at room temperature for 2 hrs. To the reaction mixture 5% NaHCO₃ (25mL) solution was added and extracted with ethyl acetate (2x25mL). Ethyl acetate layer was dried over anhydrous Na₂SO₄ and evaporated to dryness. The residue obtained was purified by flash column chromatography and eluted with 5% MeOH in CH₂Cl₂ to get 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam (14.6g, 80%).

15

2'-O-(dimethylaminoxyethyl)-5-methyluridine

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and triethylamine (1.67mL, 12mmol, dry, kept over KOH). This mixture of triethylamine-2HF was then added to 5'-O-tert-butylidiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol) and stirred at room temperature for 24 hrs. Reaction was monitored by TLC (5% MeOH in CH₂Cl₂). Solvent was removed under vacuum and the residue placed on a flash column and eluted with 10% MeOH in CH₂Cl₂ to get 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766mg, 92.5%).

25

5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine

2'-O-(dimethylaminoxyethyl)-5-methyluridine (750mg, 2.17mmol) was dried over P₂O₅ under high vacuum overnight at 40°C. It was then co-evaporated with anhydrous pyridine (20mL). The residue obtained was dissolved in pyridine (11mL) under argon atmosphere. 4-dimethylaminopyridine

30

-62-

(26.5mg, 2.60mmol), 4,4'-dimethoxytrityl chloride (880mg, 2.60mmol) was added to the mixture and the reaction mixture was stirred at room temperature until all of the starting material disappeared. Pyridine was removed under vacuum and the residue chromatographed and eluted with 10% MeOH in CH₂Cl₂ (containing a few drops of pyridine) to get 5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine (1.13g, 80%).

5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

5'-O-DMT-2'-O-(dimethylaminooxyethyl)-5-methyluridine (1.08g, 1.67mmol) was co-evaporated with toluene (20mL). To the residue N,N-diisopropylamine tetrazonide (0.29g, 1.67mmol) was added and dried over P₂O₅ under high vacuum overnight at 40°C. Then the reaction mixture was dissolved in anhydrous acetonitrile (8.4mL) and 2-cyanoethyl-N,N,N¹,N¹-tetraisopropylphosphoramidite (2.12mL, 6.08mmol) was added. The reaction mixture was stirred at ambient temperature for 4 hrs under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:ethyl acetate 1:1). The solvent was evaporated, then the residue was dissolved in ethyl acetate (70mL) and washed with 5% aqueous NaHCO₃ (40mL). Ethyl acetate layer was dried over anhydrous Na₂SO₄ and concentrated. Residue obtained was chromatographed (ethyl acetate as eluent) to get 5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%).

2'-(Aminooxyethoxy) nucleoside amidites

2'-(Aminooxyethoxy) nucleoside amidites [also known in the art as 2'-O-(aminooxyethyl) nucleoside amidites] are prepared as described in the following paragraphs.

Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

5 N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-
[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]

The 2'-O-aminooxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P. D., Guinasso, C. J., WO 94/02501 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As before the hydroxyl group may be displaced by N-hydroxyphthalimide via a Mitsunobu reaction, and the protected nucleoside may phosphitylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite].

30 2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites

2'-dimethylaminoethoxyethoxy nucleoside amidites (also known in the art as 2'-O-dimethylaminoethoxyethyl, i.e., 2'-O-CH₂-O-CH₂-N(CH₂)₂, or 2'-DMAEOE nucleoside amidites) are

-64-

prepared as follows. Other nucleoside amidites are prepared similarly.

2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine

5 2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) is slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. Hydrogen gas evolves as the solid dissolves. O²-, 2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium
10 bicarbonate (2.5 mg) are added and the bomb is sealed, placed in an oil bath and heated to 155°C for 26 hours. The bomb is cooled to room temperature and opened. The crude solution is concentrated and the residue partitioned between water (200 mL) and hexanes (200 mL). The excess
15 phenol is extracted into the hexane layer. The aqueous layer is extracted with ethyl acetate (3x200 mL) and the combined organic layers are washed once with water, dried over anhydrous sodium sulfate and concentrated. The residue is columned on silica gel using methanol/methylene
20 chloride 1:20 (which has 2% triethylamine) as the eluent. As the column fractions are concentrated a colorless solid forms which is collected to give the title compound as a white solid.

25 **5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine in anhydrous pyridine (8 mL), triethylamine (0.36 mL) and dimethoxytrityl chloride
30 (DMT-Cl, 0.87 g, 2 eq.) are added and stirred for 1 hour. The reaction mixture is poured into water (200 mL) and extracted with CH₂Cl₂ (2x200 mL). The combined CH₂Cl₂ layers are washed with saturated NaHCO₃ solution, followed by saturated NaCl solution and dried over anhydrous sodium

-65-

sulfate. Evaporation of the solvent followed by silica gel chromatography using MeOH:CH₂Cl₂:Et₃N (20:1, v/v, with 1% triethylamine) gives the title compound.

- 5 5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite
- Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) are added
- 10 to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH₂Cl₂ (20 mL) under an atmosphere of argon. The reaction mixture is stirred overnight and the solvent evaporated. The resulting residue is purified by
- 15 silica gel flash column chromatography with ethyl acetate as the eluent to give the title compound.

Example 2

Oligonucleotide synthesis

- 20 Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine.
- 25 Phosphorothioates (P=S) are synthesized as for the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the
- 30 capping step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 h), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution. Phosphinate oligonucleotides are prepared as

-66-

described in U.S. Patent 5,508,270, herein incorporated by reference.

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by
5 reference.

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as
10 described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499,
15 respectively), herein incorporated by reference.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as
20 described in U.S. Patent 5,023,243, herein incorporated by reference.

Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both
25 herein incorporated by reference.

Example 3

Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedi-
30 methylhydrazo linked oligonucleosides, also identified as MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleo-

-67-

sides, as well as mixed backbone compounds having, for instance, alternating MMI and P=O or P=S linkages are prepared as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677, 5,602,240 and 5,610,289, all of which are
5 herein incorporated by reference.

Formacetal and thioformacetal linked oligonucleosides are prepared as described in U.S. Patents 5,264,562 and 5,264,564, herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as
10 described in U.S. Patent 5,223,618, herein incorporated by reference.

Example 4

PNA Synthesis

15 Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23. They may also be prepared in accordance
20 with U.S. Patents 5,539,082, 5,700,922, and 5,719,262, herein incorporated by reference.

Example 5

Synthesis of Chimeric Oligonucleotides

25 Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between 5' and 3' "wing" segments of linked
30 nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are

-68-

also known in the art as "hemimers" or "wingmers".

[2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric

Phosphorothioate Oligonucleotides

5 Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 380B, as above. Oligonucleotides are synthesized using the automated
10 synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by increasing the wait step after the delivery of tetrazole and base to 600 s
15 repeated four times for RNA and twice for 2'-O-methyl. The fully protected oligonucleotide is cleaved from the support and the phosphate group is deprotected in 3:1 ammonia/ethanol at room temperature overnight then lyophilized to dryness. Treatment in methanolic ammonia
20 for 24 hrs at room temperature is then done to deprotect all bases and sample was again lyophilized to dryness. The pellet is resuspended in 1M TBAF in THF for 24 hrs at room temperature to deprotect the 2' positions. The reaction is then quenched with 1M TEAA and the sample is then reduced
25 to 1/2 volume by rotovac before being desalted on a G25 size exclusion column. The oligo recovered is then analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

30 [2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(Methoxyethyl)] Chimeric Phosphorothioate Oligonucleotides

[2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were

prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-methyl amidites.

- 5 [2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy
Phosphorothioate]--[2'-O-(2-Methoxyethyl)
Phosphodiester] Chimeric Oligonucleotides
[2'-O-(2-methoxyethyl phosphodiester)--[2'-deoxy phos-
phorothioate]--[2'-O-(methoxyethyl) phosphodiester]
10 chimeric oligonucleotides are prepared as per the above
procedure for the 2'-O-methyl chimeric oligonucleotide with
the substitution of 2'-O-(methoxyethyl) amidites for the
2'-O-methyl amidites, oxidization with iodine to generate
the phosphodiester internucleotide linkages within the wing
15 portions of the chimeric structures and sulfurization
utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1 dioxide (Beaucage
Reagent) to generate the phosphorothioate internucleotide
linkages for the center gap.

- Other chimeric oligonucleotides, chimeric oligonucleo-
20 sides and mixed chimeric oligonucleotides/oligonucleosides
are synthesized according to United States patent
5,623,065, herein incorporated by reference.

Example 6

25 Oligonucleotide Isolation

- After cleavage from the controlled pore glass column
(Applied Biosystems) and deblocking in concentrated
ammonium hydroxide at 55°C for 18 hours, the
oligonucleotides or oligonucleosides are purified by
30 precipitation twice out of 0.5 M NaCl with 2.5 volumes
ethanol. Synthesized oligonucleotides were analyzed by
polyacrylamide gel electrophoresis on denaturing gels and
~~judged to be at least 85% full length material.~~ The
relative amounts of phosphorothioate and phosphodiester

-70-

linkages obtained in synthesis were periodically checked by ³¹P nuclear magnetic resonance spectroscopy, and for some studies oligonucleotides were purified by HPLC, as described by Chiang et al., *J. Biol. Chem.* 1991, 266, 18162-18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

Example 7

10 Oligonucleotide Synthesis - 96 Well Plate Format

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a standard 96 well format.

15 Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1-dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster City, CA, or Pharmacia, Piscataway, NJ).

Non-standard nucleosides are synthesized as per known literature or patented methods. They are utilized as base-protected beta-cyanoethyl-diisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and deprotected with concentrated NH₄OH at elevated temperature (55-60°C) for 12-16 hours and the released product then dried in vacuo. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

-71-

Example 8**Oligonucleotide Analysis - 96 Well Plate Format**

The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

Example 9**Cell culture and oligonucleotide treatment**

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following four cell types are provided for illustrative purposes, but other cell types can be routinely used.

T-24 cells:

The transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100

-72-

micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

10

A549 cells:

The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells were routinely cultured in DMEM basal media (Gibco/Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum (Gibco/Life Technologies, Gaithersburg, MD), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Gibco/Life Technologies, Gaithersburg, MD). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

15

20

NHDF cells:

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville MD). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville MD) supplemented as recommended by the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

25

30

HEK cells:

Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville MD). HEKs were routinely maintained in ~~Keratinocyte Growth Medium~~

-73-

(Clonetics Corporation, Walkersville MD) formulated as recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

5

Treatment with antisense compounds:

When cells reached 80% confluency, they were treated with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 200 μ L OPTI-MEM™-1 reduced-serum medium (Gibco BRL) and then treated with 130 μ L of OPTI-MEM™-1 containing 3.75 μ g/mL LIPOFECTIN™ (Gibco BRL) and the desired oligonucleotide at a final concentration of 150 nM. After 4 hours of treatment, the medium was replaced with fresh medium. Cells were harvested 16 hours after oligonucleotide treatment.

15

Example 10

Analysis of oligonucleotide inhibition of liver glycogen phosphorylase expression

Antisense modulation of liver glycogen phosphorylase expression can be assayed in a variety of ways known in the art. For example, liver glycogen phosphorylase mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.2.1-4.2.9, John Wiley &

20

25

30

-74-

Sons, Inc., 1996. Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster City, CA and used according to manufacturer's instructions. Other methods of PCR are also known in the art.

Liver glycogen phosphorylase protein levels can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), ELISA or fluorescence-activated cell sorting (FACS). Antibodies directed to liver glycogen phosphorylase can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, MI), or can be prepared via conventional antibody generation methods. Methods for preparation of polyclonal antisera are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997. Preparation of monoclonal antibodies is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons, Inc., 1997.

Immunoprecipitation methods are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-10.16.11, John Wiley & Sons, Inc., 1998. Western blot (immunoblot) analysis is standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley & Sons, Inc., 1997. Enzyme-linked immunosorbent assays (ELISA) are standard in the art and can be found at, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 2, pp. 11.2.1-

11.2.22, John Wiley & Sons, Inc., 1991.

Example 11

Poly(A)+ mRNA isolation

5 Poly(A)+ mRNA was isolated according to Miura et al.,
Clin. Chem., 1996, 42, 1758-1764. Other methods for
poly(A)+ mRNA isolation are taught in, for example,
Ausubel, F.M. et al., *Current Protocols in Molecular*
Biology, Volume 1, pp. 4.5.1-4.5.3, John Wiley & Sons,
10 Inc., 1993. Briefly, for cells grown on 96-well plates,
growth medium was removed from the cells and each well was
washed with 200 μ L cold PBS. 60 μ L lysis buffer (10 mM Tris-
HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM
vanadyl-ribonucleoside complex) was added to each well, the
15 plate was gently agitated and then incubated at room
temperature for five minutes. 55 μ L of lysate was
transferred to Oligo d(T) coated 96-well plates (AGCT Inc.,
Irvine CA). Plates were incubated for 60 minutes at room
temperature, washed 3 times with 200 μ L of wash buffer (10
20 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the
final wash, the plate was blotted on paper towels to remove
excess wash buffer and then air-dried for 5 minutes. 60 μ L
of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C
was added to each well, the plate was incubated on a 90°C
25 hot plate for 5 minutes, and the eluate was then
transferred to a fresh 96-well plate.

Cells grown on 100 mm or other standard plates may be
treated similarly, using appropriate volumes of all
solutions.

-76-

Example 12**Total RNA Isolation**

Total mRNA was isolated using an RNEASY 96™ kit and buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200 µL cold PBS. 100 µL Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 100 µL of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96™ well plate attached to a QIAVAC™ manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1 mL of Buffer RW1 was added to each well of the RNEASY 96™ plate and the vacuum again applied for 15 seconds. 1 mL of Buffer RPE was then added to each well of the RNEASY 96™ plate and the vacuum applied for a period of 15 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVAC™ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC™ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 60 µL water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60 µL water.

The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are

-77-

carried out.

Example 13

Real-time Quantitative PCR Analysis of liver glycogen 5 phosphorylase mRNA Levels

Quantitation of liver glycogen phosphorylase mRNA levels was determined by real-time quantitative PCR using the ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's
10 instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time. As opposed to standard PCR, in which amplification products are quantitated after the PCR is
15 completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye
20 (e.g., JOE or FAM, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City, CA) is attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, obtained from either Operon Technologies Inc., Alameda, CA or PE-Applied Biosystems, Foster City,
25 CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-
30 exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is

-78-

generated. With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

10 PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25 µL PCR cocktail (1x TAQMAN™ buffer A, 5.5 mM MgCl₂, 300 µM each of dATP, dCTP and dGTP, 600 µM of dUTP, 100 nM each of forward primer, reverse primer, and probe, 15 20 Units RNase inhibitor, 1.25 Units AMPLITAQ GOLD™, and 12.5 Units MuLV reverse transcriptase) to 96 well plates containing 25 µL poly(A) mRNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C to activate the 20 AMPLITAQ GOLD™, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension). Liver glycogen phosphorylase probes and primers were designed to hybridize to the human liver glycogen phosphorylase sequence, using 25 published sequence information (GenBank accession number M14636, incorporated herein as SEQ ID NO:1).

For liver glycogen phosphorylase the PCR primers were: forward primer: CATGGGCCGAACATTACAGAA (SEQ ID NO: 2) reverse primer: CAAGACCACCATTTGCCAAGTC (SEQ ID NO: 3) and 30 the PCR probe was: FAM-CTGTGATGAGGCCATTTACCAGCTTGG-TAMRA (SEQ ID NO: 4) where FAM (PE-Applied Biosystems, Foster City, CA) is the ~~Fluorescent reporter dye~~ and TAMRA (PE-

-79-

Applied Biosystems, Foster City, CA) is the quencher dye.

For GAPDH the PCR primers were:

forward primer: GAAGGTGAAGGTCGGAGTC (SEQ ID NO: 5)

reverse primer: GAAGATGGTGATGGGATTTC (SEQ ID NO: 6) and the

5 PCR probe was: 5' JOE-CAAGCTTCCCGTTCTCAGCC- TAMRA 3' (SEQ ID NO: 7) where JOE (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

10 Example 14

Northern blot analysis of liver glycogen phosphorylase mRNA levels

Eighteen hours after antisense treatment, cell monolayers were washed twice with cold PBS and lysed in 1 mL RNAZOL™ (TEL-TEST "B" Inc., Friendswood, TX). Total RNA was prepared following manufacturer's recommended protocols. Twenty micrograms of total RNA was fractionated by electrophoresis through 1.2% agarose gels containing 1.1% formaldehyde using a MOPS buffer system (AMRESCO, Inc. Solon, OH). RNA was transferred from the gel to HYBOND™-N+ nylon membranes (Amersham Pharmacia Biotech, Piscataway, NJ) by overnight capillary transfer using a Northern/Southern Transfer buffer system (TEL-TEST "B" Inc., Friendswood, TX). RNA transfer was confirmed by UV visualization. Membranes were fixed by UV cross-linking using a STRATALINKER™ UV Crosslinker 2400 (Stratagene, Inc, La Jolla, CA).

Membranes were probed using QUICKHYB™ hybridization solution (Stratagene, La Jolla, CA) using manufacturer's recommendations for stringent conditions with a liver glycogen phosphorylase specific probe prepared by PCR using the forward primer CATGGGCCGAACATTACAGAA (SEQ ID NO: 2) and the reverse primer CAAGACCACCATTTGCCAAGTC (SEQ ID NO: 3).

-80-

To normalize for variations in loading and transfer efficiency membranes were stripped and probed for glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA). Hybridized membranes were
5 visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3 (Molecular Dynamics, Sunnyvale, CA). Data was normalized to GAPDH levels in untreated controls.

10 Example 15

Antisense inhibition of liver glycogen phosphorylase expression- phosphorothioate oligodeoxynucleotides

In accordance with the present invention, a series of oligonucleotides were designed to target different regions
15 of the human liver glycogen phosphorylase RNA, using published sequences (GenBank accession number M14636, incorporated herein as SEQ ID NO: 1). The oligonucleotides are shown in Table 1. Target sites are indicated by nucleotide numbers, as given in the sequence source
20 reference (Genbank accession no. M14636), to which the oligonucleotide binds. All compounds in Table 1 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. The compounds were analyzed for effect on liver glycogen phosphorylase mRNA
25 levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

-81-

Table 1

Inhibition of liver glycogen phosphorylase mRNA levels by
phosphorothioate oligodeoxynucleotides

ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
104049	5' UTR	9	Ccgccccgcgcgccaggag	0	8
104050	5' UTR	71	Cgggctgcgcagagagctgg	13	9
104051	Start Codon	109	Cagcggttcgcccatggctg	2	10
104052	Start Codon	114	Tctgtcagcggttcgcccat	31	11
104053	Coding	172	Tgccacgttctccacgccca	82	12
104054	Coding	220	Cttgaccagcgtgaagtgca	52	13
104055	Coding	260	Gcgcgaagtagtagtcgcgg	20	14
104056	Coding	299	Tccagcgccccaccaggtgg	56	15
104057	Coding	372	Cccatgtaaaattccagaga	56	16
104058	Coding	415	Attttgcagaccgaggttga	69	17
104059	Coding	460	Ctcttctatatccaatccaa	37	18
104060	Coding	523	Gcaggcagcaagtctcccaa	71	19
104061	Coding	571	Gccgtatccataggctgcaa	0	20
104062	Coding	625	Tacctgccatccatctcgga	85	21
104063	Coding	678	Gggcgggacttctcccaagg	21	22
104064	Coding	734	Tcccgggtgttggtgtgttct	37	23
104065	Coding	817	Ggtgttgacagtgttattca	44	24
104066	Coding	877	Tccaacattaaagtctctga	38	25
104067	Coding	949	Attgtcattgggatagagga	51	26
104068	Coding	1052	Tggagccaaacttggaggct	68	27
104069	Coding	1206	Ttctgggtgagctcccatgc	48	28
104070	Coding	1263	Acgggcccagcgctccagggc	53	29
104071	Coding	1327	Atgcttctgatttatctcat	51	30
104072	Coding	1389	Atcagagacatccttctcag	67	31
104073	Coding	1453	Cacagcatgggaaccgacaa	85	32
104074	Coding	1510	Gaagtccttgaatactttag	60	33
104075	Coding	1598	Ctgcaagtcctgggttgca	51	34
104076	Coding	1678	Atcaccaggaagctgtgga	69	35
104077	Coding	1778	Aggatgggttgatcttcact	61	36
104078	Coding	1837	Acagttcaagagctgtcgct	61	37
104079	Coding	1925	Cagctttaccaccaatgata	71	38
104080	Coding	2009	Ttccaaccatagggtcattg	73	39
104081	Coding	2080	Atctgtggctggaatgactt	21	40
104082	Coding	2135	Tcatattgcctgtccccgag	82	41
104083	Coding	2189	Ccacattggcccatccatg	81	42
104084	Coding	2241	Atgctcatgccaaagatgaa	2	43

-82-

104085	Coding	2291	Agtattcttttgcctcgtac	79	44
104086	Coding	2337	Ttgtcaatttgatcaatgac	38	45
104087	Coding	2403	Tcatgataaaatagcatgtt	0	46
104088	3' UTR	2758	Cccattcccagagatactc	68	47

As shown in Table 1, SEQ ID NOS 11, 12, 13, 15, 16, 17, 18, 19, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 44, 45 and 47 demonstrated at least 30% inhibition of liver glycogen phosphorylase expression in this assay and are therefore preferred.

Example 16:

Antisense inhibition of liver glycogen phosphorylase

10 expression- phosphorothioate 2'-MOE gapmer oligonucleotides

In accordance with the present invention, a second series of oligonucleotides targeted to human liver glycogen phosphorylase were synthesized. The oligonucleotide sequences are shown in Table 2. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. M14636), to which the oligonucleotide binds.

All compounds in Table 2 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. Cytidine residues in the 2'-MOE wings are 5-methylcytidines.

Data were obtained by real-time quantitative PCR as described in other examples herein and are averaged from two experiments. If present, "N.D." indicates "no data".

-83-

Table 2

Inhibition of liver glycogen phosphorylase mRNA levels by
chimeric phosphorothioate oligonucleotides having 2'-MOE
wings and a deoxy gap

5

ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
104089	5' UTR	9	Ccgccccgcccgcgccaggag	0	8
104090	5' UTR	71	Cgggctgcgccagagagctgg	61	9
104091	Start Codon	109	Cagcgggttcgcccattggctg	20	10
104092	Start Codon	114	Tctgtcagcgggttcgcccatt	23	11
104093	Coding	172	Tgccacgttctccacgccc	90	12
104094	Coding	220	Cttgaccagcgtgaagtgc	88	13
104095	Coding	260	Gcgcgaaagtagtagtcgcg	76	14
104096	Coding	299	Tccagcgcgccaccagggtg	87	15
104097	Coding	372	Cccatgtaaaattccagaga	98	16
104098	Coding	415	Attttgcagaccgaggttga	94	17
104099	Coding	460	Ctcttctatatccaatccaa	71	18
104100	Coding	523	Gcaggcagcaagtctcccaa	57	19
104101	Coding	571	Gccgtatccataggctgcaa	97	20
104102	Coding	625	Tacctgccatccatctcgga	89	21
104103	Coding	678	Gggcgaggacttctcccaagg	79	22
104104	Coding	734	Tcccgggtgttggtgtgttct	89	23
104105	Coding	817	Ggtgttgacagtgttattca	95	24
104106	Coding	877	Tccaacattaaagtctctga	77	25
104107	Coding	949	Attgtcattgggatagagga	92	26
104108	Coding	1052	Tggagccaaacttggaggct	77	27
104109	Coding	1206	Ttctgggttgagctcccatgc	48	28
104110	Coding	1263	Acggggccagcgtccagggc	82	29
104111	Coding	1327	Atgcttctgatttatctcat	83	30
104112	Coding	1389	Atcagagacatccttctcag	96	31
104113	Coding	1453	Cacagcatgggaaccgacaa	88	32
104114	Coding	1510	Gaagtccttgaatactttag	86	33
104115	Coding	1598	Ctgcaagtccctgggttgca	74	34
104116	Coding	1678	Atcaccaggaagctgtgga	70	35
104117	Coding	1778	Aggatgggttgatcttcact	70	36
104118	Coding	1837	Acagttcaagagctgtcgct	85	37
104119	Coding	1925	Cagctttaccaccaatgata	58	38
104120	Coding	2009	Ttccaaccataggggtcattg	86	39
104121	Coding	2080	Atctgtggctggaatgactt	40	40
104122	Coding	2135	Tcatattgctgtccccgag	49	41

-84-

104123	Coding	2189	Ccacattggcccccattccatg	90	42
104124	Coding	2241	Atgctcatgccaaagatgaa	53	43
104125	Coding	2291	Agtattcttttgctcgtac	86	44
104126	Coding	2337	Ttgtcaatttgatcaatgac	73	45
104127	Coding	2403	Tcatgataaaaatagcatgtt	71	46
104128	3' UTR	2758	Ccccattcccagagatactc	82	47

As shown in Table 2, SEQ ID NOS 9, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46 and 47 demonstrated at least 30% inhibition of liver glycogen phosphorylase expression in this experiment and are therefore preferred.

Example 17

10 Western blot analysis of liver glycogen phosphorylase protein levels

Western blot analysis (immunoblot analysis) is carried out using standard methods. Cells are harvested 16-20 h after oligonucleotide treatment, washed once with PBS, 15 suspended in Laemmli buffer (100 ul/well), boiled for 5 minutes and loaded on a 16% SDS-PAGE gel. Gels are run for 1.5 hours at 150 V, and transferred to membrane for western blotting. Appropriate primary antibody directed to liver glycogen phosphorylase is used, with a radiolabelled or 20 fluorescently labeled secondary antibody directed against the primary antibody species. Bands are visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA).

Example 18

25 Cell culture of primary mouse hepatocytes

Primary mouse hepatocytes were prepared from CD-1 mice purchased from Charles River Labs (Wilmington, MA) and were routinely cultured in Hepatocyte Attachment Media (Gibco) supplemented with 10% Fetal Bovine Serum (Gibco/Life

-85-

Technologies, Gaithersburg, MD), 250nM dexamethasone (Sigma), 10nM bovine insulin (Sigma). Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 10000 cells/well for use in RT-PCR analysis.

5 For Northern blotting or other analyses, cells may be seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide. In Example 19 below, cells are treated with 150 nM oligonucleotide.

10

Example 19

Antisense inhibition of mouse liver glycogen phosphorylase expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap.

15 In accordance with the present invention, a third series of oligonucleotides were designed to target different regions of mouse liver glycogen phosphorylase RNA, using a consensus sequence incorporated herein as SEQ ID NO: 48 which was prepared from alignment of the
20 following 21 published sequences (GenBank accession numbers D17979.1, W65769.1, AA049580.1, AA208939.1, AA212447.1, AA246021.1, AA427221.1, AA792928.1, AI118771.1, AI180947.1, AI194480.1, AI196311.1, AI256150.1, AI303555.1, AI507050.1, AI507164.1, AI605246.1, AA492865.1, AA511651.1, 25 AI615434.1, and AI597375.1). The oligonucleotides are shown in Table 3. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 3 are chimeric oligonucleotides ("gapmers") 20 nucleotides in
30 length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate

-86-

(P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on mouse liver glycogen phosphorylase mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments. If present, "N.D." indicates "no data".

Table 3

Inhibition of mouse liver glycogen phosphorylase mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO
115885	start codon	48	34	Tttgccatgggttgcgggcgg	76	49
115886	coding	48	64	Tgccgtcgcttctcctggtc	96	50
115887	coding	48	80	Gcctcggatgctgatctgcc	95	51
115888	coding	48	101	Cacattctctacgcccacga	92	52
115889	coding	48	124	Ttgaaaccctttttcagctc	93	53
115890	coding	48	151	Ttgaccagagtgaagtgcag	90	54
115891	coding	48	253	Gtagtagtgctgctgtgtac	84	55
115892	coding	48	262	Acacttgctcgtagtagtgct	74	56
115893	coding	48	278	Aatacaccctcttgggacac	82	57
115894	coding	48	287	Gagagaggtaatacaccctc	86	58
115895	coding	48	300	Atgtaaaattccagagagag	75	59
115896	coding	48	301	Catgtaaaattccagagaga	75	60
115897	coding	48	343	Ttgtaagccaagggtgatca	97	61
115898	coding	48	347	Catttttgaagccaagggttg	79	62
115899	coding	48	380	Ccaatccaagctggtaaatg	86	63
115900	coding	48	400	Gtcttctaactcttccatgt	92	64
115901	coding	48	423	Gcaaggccggcatcttcttc	75	65
115902	coding	48	428	Cattcgcaaggccggcatct	70	66
115903	coding	48	505	Gaatgccgtacgatagggct	0	67
115904	coding	48	520	Taccgtattcataacgaatg	55	68
115905	coding	48	662	Ttgtcaccgctgggtgtgtc	0	69
115906	genomic	48	749	Ggtcccaggtgccatgaagt	47	70
115907	genomic	48	754	Ctgcaggtcccaggtgccat	0	71
115908	genomic	48	768	Aaattcgtagatagctgcag	7	72
115909	genomic	48	797	Tctgctgggtgtgtggttaagt	51	73
115910	genomic	48	825	Gaccgcgagcacagccaccg	52	74
115911	genomic	48	830	Cccgagaccgcgagcacagc	59	75
115912	coding	48	890	Tggctcaggtctttcacata	89	76
115913	coding	48	918	Actcacaactgtggagct	91	77
115914	coding	48	934	Ggaggaagatgtcatcctc	81	78

115915	coding	48	986	Aggaactgggagaatttcag	65	79
115916	coding	48	990	Ctccaggaactgggagaatt	74	80
115917	coding	48	1005	Caccttgattccttctcca	92	81
115918	coding	48	1057	Actcgtggatccgcttcacg	98	82
115919	coding	48	1068	Ctgccttttatactcgtgga	90	83
115920	coding	48	1087	Catgcaggcagttcagaagc	31	84
115921	coding	48	1100	Tacatggtgatcacatgcag	70	85
115922	coding	48	1138	Gcatgaacttcatgttgctt	94	86
115923	coding	48	1178	Ttggcccatccatagtccc	89	87
115924	genomic	48	1284	Atattctttggcctcatacc	74	88
115925	genomic	48	1287	Ctaatattctttggcctcat	84	89
115926	genomic	48	1300	Agcagccaatcatctaata	43	90
115927	genomic	48	1307	Ttatccaagcagccaatcat	12	91
115928	genomic	48	1337	Ggcctcgtatattcttggc	5	92
115929	genomic	48	1340	Gagggccctcgtatattctt	0	93
115930	genomic	48	1362	Tcgtatgaccaaactcagttc	0	94
115931	genomic	48	1363	Gtcgtatgaccaaactcagtt	56	95
115932	coding	48	1400	Tctggctgattgggagaaaa	91	96
115933	coding	48	1401	Gtctggctgattgggagaaa	94	97
115934	coding	48	1457	Gcaaagactttaaatctgtc	88	98
115935	coding	48	1479	Cttgacataggcttcgtagt	86	99
115936	coding	48	1487	Tcttgacacttgacataggc	85	100
115937	coding	48	1492	Ctttttcttgacacttgaca	76	101
115938	coding	48	1504	Acagctgactgactttttct	85	102
115939	coding	48	1519	Ctttttgattcatatacagc	76	103
115940	coding	48	1529	Gtgttccaggctttttgatt	87	104
115941	coding	48	1567	Agaacttccctgaggcagct	95	105
115942	coding	48	1572	Actggagaacttccctgagg	85	106
115943	coding	48	1591	Actccttaattgttcggtca	96	107
115944	coding	48	1656	Gctggactcgttggatagg	99	108
115945	coding	48	1661	Ccattgctggactcgttgg	99	109
115946	coding	48	1668	Actgacccattgctggact	97	110
115947	coding	48	1673	Ttggcactgacccattgct	94	111
115948	coding	48	1676	Ccattggcactgacccatt	96	112
115949	stop codon	48	1695	Acatttttagcattcacttcc	94	113
115950	3'UTR	48	1738	Ctctaaaacactcaagttcc	86	114
115951	genomic	48	1747	Tcaggggaagctctaaaacac	89	115
115952	genomic	48	1792	Ttcccatccccggagacact	98	116
115953	genomic	48	1907	Tgagagctctaggaaatcct	46	117
115954	genomic	48	1911	Tccctgagagctctaggaaa	30	118
115955	genomic	48	1913	Agtccctgagagctctagg	30	119
115956	genomic	48	1933	Tactcttcggttggtggttt	42	120
115957	genomic	48	1961	Tggctggagtcatgggtccc	44	121
115958	genomic	48	1970	Ctctttatgtggctggagtc	71	122
115959	genomic	48	2010	Ccaagggcctcccctccac	68	123
115960	genomic	48	2053	Gctctagcatccccctacac	38	124
115961	genomic	48	2066	Ctgcctcatcatggctctag	33	125
115962	genomic	48	2067	Cctgcctcatcatggctcta	54	126

-88-

As shown in Table 3, SEQ ID NOs 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 75, 76, 77, 78, 79, 80, 81, 82, 83, 85, 86, 87, 88, 89, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 122 and 123 demonstrated at least 59% inhibition of mouse liver glycogen phosphorylase expression in this experiment and are therefore preferred. The target sites to which these preferred sequences are complementary are herein referred to as "active sites" and are therefore preferred sites for targeting by compounds of the present invention.

Example 20

Antisense inhibition of mouse liver glycogen phosphorylase expression (ISIS 115890): dose response

ISIS 115890 (SEQ ID NO: 54) was identified as an oligonucleotide with good activity based on the results in Example 19 and was further investigated in a dose response experiment. Mouse hepatocytes were dosed with 5, 25, 100 and 250 nM of ISIS 115890 and exhibited reduction in mRNA expression of 2%, 42%, 77% and 83% respectively. These results indicate that reduction of mouse liver glycogen phosphorylase expression by ISIS 115890 is dependent on the oligonucleotide dosage.

25

Example 21

Effects of antisense inhibition of mouse liver glycogen phosphorylase (ISIS 115922 and ISIS 115890) on blood glucose levels

db/db mice are used as a model of Type 2 diabetes. These mice are hyperglycemic, obese, hyperlipidemic, and insulin resistant. The db/db phenotype is due to a ~~mutation in the leptin receptor~~ on a C57BLKS background. However, a mutation in the leptin gene on a different mouse

-89-

background can produce obesity without diabetes (ob/ob mice). Leptin is a hormone produced by fat that regulates appetite and animals or humans with leptin deficiencies become obese. Heterozygous db/wt mice (known as lean littermates) do not display the hyperglycemia/hyperlipidemia or obesity phenotype.

In accordance with the present invention, ISIS 115922 (GCATGAACCTTCATGTTGCCT, SEQ ID No: 86) and ISIS 115890 (TTGACCAGAGTGAAGTGCAG, SEQ ID NO: 54) were investigated in experiments designed to address the role of liver glycogen phosphorylase in glucose metabolism and homeostasis. ISIS 115922 is completely complementary to sequences of the mouse liver glycogen phosphorylase nucleotide sequence incorporated herein as SEQ ID NO: 48 (starting at nucleotide 1138). ISIS 115890 is completely complementary to sequences of the mouse liver glycogen phosphorylase nucleotide sequence incorporated herein as SEQ ID NO: 48 (starting at nucleotide 151). The control used is ISIS 29848 (NNNNNNNNNNNNNNNNNNNNNN, SEQ ID NO: 127) where N is a mixture of A, G, T and C.

8-week old male db/db mice (n=8/group) were dosed twice weekly (intraperitoneal injection) for seven weeks with saline, ISIS 115922 at 10, 25, 50 and 75 mg/kg/week, ISIS 115890 at 50 mg/kg/week or ISIS 29848 (universal control) at 50 mg/kg/week. Blood glucose levels (in the fed state) were measured weekly.

By week 6, ISIS 115890 at 50 mg/kg/week lowered the glucose levels 100 mg/dL relative to the saline-treated control and 65 mg/dL relative to the ISIS 29848 whereas ISIS 115922 at 50 mg/kg/week had a more modest effect, lowering glucose levels by 35 mg/kg relative to the saline-treated control and 16 mg/dL relative to ISIS 29848.

After 4 weeks of treatment, a glucose tolerance test was conducted in 16-hour fasted mice. Glucose levels were

-90-

measured 30, 60 and 120 min. following administration of glucose. Statistically, there was no improved glucose tolerance in mice treated with the antisense oligonucleotides relative to mice treated with saline or
5 ISIS 29848.

In a similar experiment, ob/ob mice (n=8/group) were dosed twice weekly (intraperitoneal injection) for 5 weeks with saline or ISIS 115890 at 25 mg/kg per dose. Blood glucose was measured every week and an oral glucose
10 tolerance test (OGTT) was performed 3 weeks after initiation of treatment.

By week 5, ISIS 115890 lowered the glucose levels by 76 mg/dL relative to the saline-treated ob/ob mice.

After 3 weeks of treatment, the OGTT was conducted in
15 16 hour fasted mice. Glucose levels were measured 30, 60 and 120 minutes following administration of glucose. ob/ob mice treated with ISIS 115890 exhibited glucose levels 50-75 mg/dL lower than the saline-treated ob/ob mice, indicating an improved glucose tolerance for animals
20 treated with ISIS 115890.

Example 22

Antisense inhibition of mouse liver glycogen phosphorylase expression (ISIS 115922): dose response

25 ISIS 115922 (SEQ ID NO: 86) was identified as an oligonucleotide with good activity based on the results in Example 19 and was further investigated in a dose response experiment. Following sacrifice of the db/db mice after the 6-week investigation described in Example 21, liver samples
30 were collected and liver glycogen phosphorylase mRNA levels were determined by RT-PCR (N=4/group). Mice dosed with 10, 25, 50 and 75 mg of ISIS 115922 exhibited reduction in mRNA expression of 0%, 25%, 50% and 75% respectively, relative to saline treated mice. These results indicate that

-91-

reduction of mouse liver glycogen phosphorylase by ISIS 115922 is dependent on the oligonucleotide dosage.

Example 23

- 5 Effects of antisense inhibition of mouse liver glycogen phosphorylase (ISIS 115922 and ISIS 115890) on body and organ weight and levels of serum transaminases, serum lipids, serum insulin and glycogen

Following sacrifice of the db/db mice after the 6 week
10 investigation described in Example 21, serum was collected and body and organ weights were measured. No significant changes in body weights were observed. Liver weights were slightly elevated in antisense oligonucleotide treated animals, however there were no changes in fat or spleen
15 weights among the different groups.

No significant changes in serum transaminases were observed, indicating no ongoing toxic effects of the oligonucleotide treatment.

- No significant changes in serum lipid levels were
20 observed with the exception of a slight increase in triglycerides in the group dosed with 50 mg/kg/week of ISIS 115922.

- Liver samples were analyzed for glycogen content. ISIS 115922 increased hepatic glycogen content up to 2.5 fold
25 compared to the saline and universal control treated groups.

- Following sacrifice of the ob/ob mice used in the investigation described in Example 21, serum was analyzed for insulin and lipid levels. Levels of insulin were 25
30 ng/ml and 35 ng/ml lower than the saline control at weeks 4 and 5, respectively. At week 5, the animals treated with ISIS 115890 did not have cholesterol levels significantly lower than saline-treated mice. Triglyceride levels were

-92-

found to be 17% lower in animals treated with ISIS 115890 relative to the saline-treated controls.

Example 24

5 Modulation of an insulin-signaling pathway via antisense inhibition of mouse liver glycogen phosphorylase

Insulin-signaling pathways are defined as metabolic pathways regulated by insulin. Type 2 diabetes is characterized by impaired insulin-signaling pathways and increased hepatic glucose production which causes
10 hyperglycemia (Virkamäki et al., *J. Clin. Invest.* 1999, 103, 931-943). The diabetic liver can over-produce glucose via two insulin signaling pathways: increased gluconeogenesis (glucose synthesis from lactate, glycerol
15 and amino acids) and increased glycogenolysis (breakdown of glycogen, a cross-linked storage form of glucose). Thus, inhibiting insulin signaling pathways may have therapeutic potential by reducing hyperglycemia.

Phosphoenolpyruvate carboxykinase (PEPCK) is a target
20 for regulation of gluconeogenesis. Transcription of the PEPCK gene is regulated by insulin, glucocorticoids, cAMP, and diet, in order to adjust glucose production to physiologic requirements.

In accordance with the present invention, 8-week old
25 male db/db mice (n=8/group) were dosed twice weekly (intraperitoneal injection) for six weeks with saline, ISIS 115922 at 10, 25, 50 and 75 mg/kg/week, ISIS 115890 at 50 mg/kg/week or ISIS 29848 (universal control) at 50 mg/kg/week. Upon sacrifice, liver samples were collected to
30 evaluate levels of PEPCK RNA by RT-PCR (n=4/group). RNA levels were normalized and expressed as a percentage of saline treated control group.

No consistent increases in PEPCK RNA were observed, suggesting that while glycogenolysis may be inhibited, the

-93-

counteractive signal to increase glucose levels by gluconeogenesis did not occur.

-94-

What is claimed is:

1. An antisense compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding liver glycogen phosphorylase, wherein said antisense compound specifically hybridizes with and inhibits the expression of liver glycogen phosphorylase.
2. The antisense compound of claim 1 which is an antisense oligonucleotide.
3. The antisense compound of claim 2 wherein the antisense oligonucleotide has a sequence comprising SEQ ID NO: 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 75, 76, 77, 78, 79, 80, 81, 82, 83, 85, 86, 87, 88, 89, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 122 or 123.
4. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.
5. The antisense compound of claim 4 wherein the modified internucleoside linkage is a phosphorothioate linkage.
6. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.
7. The antisense compound of claim 6 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

8. The antisense compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.
9. The antisense compound of claim 8 wherein the modified nucleobase is a 5-methylcytosine.
10. The antisense compound of claim 2 wherein the antisense oligonucleotide is a chimeric oligonucleotide.
11. A pharmaceutical composition comprising the antisense compound of claim 1 and a pharmaceutically acceptable carrier or diluent.
12. The pharmaceutical composition of claim 11 further comprising a colloidal dispersion system.
13. The pharmaceutical composition of claim 11 wherein the antisense compound is an antisense oligonucleotide.
14. A compound 8 to 50 nucleobases in length which specifically hybridizes with at least an 8-nucleobase portion of an active site on a nucleic acid molecule encoding liver glycogen phosphorylase.
15. A method of inhibiting the expression of liver glycogen phosphorylase in cells or tissues comprising contacting said cells or tissues with the antisense compound of claim 1 so that expression of liver glycogen phosphorylase is inhibited.

-96-

16. The method of claim 15 wherein the cells or tissues are human cells or tissues.

17. The method of claim 15 wherein the cells or tissues are rodent cells or tissues.

18. The method of claim 17 wherein the rodent cells or tissues are mouse cells or tissues.

19. The method of claim 15 wherein the cells or tissues are liver cells or tissues.

20. A method of treating an animal having or suspected of having a disease or condition associated with liver glycogen phosphorylase comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of claim 1 so that expression of liver glycogen phosphorylase is inhibited.

21. The method of claim 20 wherein the animal is a human or a rodent.

22. The method of claim 20 wherein the disease or condition is a metabolic disease or condition.

23. The method of claim 20 wherein the disease or condition is diabetes.

24. The method of claim 20 wherein the disease or condition is Type 2 diabetes.

25. The method of claim 20 wherein the disease or condition is obesity.

26. The method of claim 20 wherein the disease or

-97-

condition is a hyperproliferative condition.

27. The method of claim 26 wherein the hyperproliferative condition is cancer.

28. A method of decreasing blood glucose levels in an animal comprising administering to said animal the compound of claim 1.

29. The method of claim 28 wherein the animal is a human or a rodent.

30. The method of claim 28 wherein the blood glucose levels are serum glucose levels.

31. The method of claim 28 wherein the animal is a diabetic animal.

32. A method of preventing or delaying the onset of a disease or condition associated with liver glycogen phosphorylase in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of claim 1.

33. The method of claim 32 wherein the animal is a human or a rodent.

34. The method of claim 32 wherein the disease or condition is a metabolic disease or condition.

35. The method of claim 32 wherein the disease or condition is diabetes.

36. The method of claim 32 wherein the disease or condition is Type 2 diabetes.

-98-

37. The method of claim 32 wherein the disease or condition is obesity.

38. The method of claim 32 wherein the disease or condition is a hyperproliferative condition.

39. The method of claim 32 wherein the hyperproliferative condition is cancer.

40. A method of preventing or delaying the onset of an increase in blood glucose levels in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of claim 1.

41. The method of claim 40 wherein the animal is a human or a rodent.

42. The method of claim 40 wherein the blood glucose levels are serum glucose levels.

43. The method of claim 40 wherein the animal is a diabetic animal.

44. A method of modulating an insulin-signaling pathway in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of an inhibitor of liver glycogen phosphorylase.

45. The method of claim 44 wherein the insulin-signaling pathway is gluconeogenesis.

-99-

46. The method of claim 44 wherein the insulin-signaling pathway is glycogenolysis.

47. The method of claim 44 wherein the animal is a human or a rodent.

48. A method of modulating an insulin-signaling pathway in an animal comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of claim 1.

49. The method of claim 48 wherein the insulin-signaling pathway is gluconeogenesis.

50. The method of claim 48 wherein the insulin-signaling pathway is glycogenolysis.

51. The method of claim 48 wherein the animal is a human or a rodent.

SEQUENCE LISTING

<110> Isis Pharmaceuticals, Inc.
 Brett P. Monia
 Lex M. Cowsert

<120> ANTISENSE MODULATION OF LIVER GLYCOGEN PHOSPHORYLASE EXPRESSION

<130> ISPH-0735

<150> US 10/019,470

<151> 2002-05-09

<150> US 10/114,544

<151> 2002-04-01

<150> PCT/US00/19019

<151> 2000-07-12

<150> US 09/357,071

<151> 1999-07-19

<160> 47

<210> 1

<211> 2828

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (114)..(2657)

<400> 1

gttgaaagct cctggcgcg cgggggcgac tccacccctg cccggcagcc cagcgctcc 60

ggcgcactt ccagctctct gcgcagcccg ccgcgcagcc cgccgcccc gcc atg 116
 Met
 1

ggc gaa ccg ctg aca gac cag gag aag cgg cgg cag atc agc atc cgc 164
 Gly Glu Pro Leu Thr Asp Gln Glu Lys Arg Arg Gln Ile Ser Ile Arg
 5 10 15

ggc atc gtg ggc gtg gag aac gtg gca gag ctg aag aag agt ttc aac 212
 Gly Ile Val Gly Val Glu Asn Val Ala Glu Leu Lys Lys Ser Phe Asn
 20 25 30

cgg cac ctg cac ttc acg ctg gtc aag gac cgc aac gtg gcc acc acc 260
 Arg His Leu His Phe Thr Leu Val Lys Asp Arg Asn Val Ala Thr Thr
 35 40 45

cgc gac tac tac ttc gcg ctg gcg cac acg gtg cgg gac cac ctg gtg 308
 Arg Asp Tyr Tyr Phe Ala Leu Ala His Thr Val Arg Asp His Leu Val
 50 55 60 65

ggg cgc tgg atc cgc acg cag cag cac tac tac gac aag tgc ccc aag 356
 Gly Arg Trp Ile Arg Thr Gln Gln His Tyr Tyr Asp Lys Cys Pro Lys
 70 75 80

agg gaa tat tac ctc tct ctg gaa ttt tac atg ggc cga aca tta cag Arg Glu Tyr Tyr Leu Ser Leu Glu Phe Tyr Met Gly Arg Thr Leu Gln 85 90 95	404
aac acc atg atc aac ctc ggt ctg caa aat gcc tgt gat gag gcc att Asn Thr Met Ile Asn Leu Gly Leu Gln Asn Ala Cys Asp Glu Ala Ile 100 105 110	452
tac cag ctt gga ttg gat ata gaa gag tta gaa gaa att gaa gaa gat Tyr Gln Leu Gly Leu Asp Ile Glu Glu Leu Glu Glu Ile Glu Glu Asp 115 120 125	500
gct gga ctt ggc aat ggt ggt ctt ggg aga ctt gct gcc tgc ttc ttg Ala Gly Leu Gly Asn Gly Gly Leu Gly Arg Leu Ala Ala Cys Phe Leu 130 135 140 145	548
gat tcc atg gca acc ctg gga ctt gca gcc tat gga tac ggc att cgg Asp Ser Met Ala Thr Leu Gly Leu Ala Ala Tyr Gly Tyr Gly Ile Arg 150 155 160	596
tat gaa tat ggg att ttc aat cag aag atc cga gat gga tgg cag gta Tyr Glu Tyr Gly Ile Phe Asn Gln Lys Ile Arg Asp Gly Trp Gln Val 165 170 175	644
gaa gaa gca gat gat tgg ctc aga tat gga aac cct tgg gag aag tcc Glu Glu Ala Asp Asp Trp Leu Arg Tyr Gly Asn Pro Trp Glu Lys Ser 180 185 190	692
cgc cca gaa ttc atg ctg cct gtg cac ttc tat gga aaa gta gaa cac Arg Pro Glu Phe Met Leu Pro Val His Phe Tyr Gly Lys Val Glu His 195 200 205	740
acc aac acc ggg acc aag tgg att gac act caa gtg gtc ctg gct ctg Thr Asn Thr Gly Thr Lys Trp Ile Asp Thr Gln Val Val Leu Ala Leu 210 215 220 225	788
cca tat gac acc ccc gag ccc ggc tac atg aat aac act gtc aac acc Pro Tyr Asp Thr Pro Glu Pro Gly Tyr Met Asn Asn Thr Val Asn Thr 230 235 240	836
atg cgc ctc tgg tct gct cgg gca cca aat gac ttt aac ctc aga gac Met Arg Leu Trp Ser Ala Arg Ala Pro Asn Asp Phe Asn Leu Arg Asp 245 250 255	884
ttt aat gtt gga gac tac att cag gct gtg ctg gac cga aac ctg gcc Phe Asn Val Gly Asp Tyr Ile Gln Ala Val Leu Asp Arg Asn Leu Ala 260 265 270	932
gag aac atc tcc cgg gtc ctc tat ccc aat gac aat ttt ttt gaa ggg Glu Asn Ile Ser Arg Val Leu Tyr Pro Asn Asp Asn Phe Phe Glu Gly 275 280 285	980
aag gag cta aga ttg aag cag gaa tac ttt gtg gtg gct gca acc ttg Lys Glu Leu Arg Leu Lys Gln Glu Tyr Phe Val Val Ala Ala Thr Leu 290 295 300 305	1028
caa gat atc atc cgc cgt ttc aaa gcc tcc aag ttt ggc tcc acc cgt Gln Asp Ile Ile Arg Arg Phe Lys Ala Ser Lys Phe Gly Ser Thr Arg	1076

3

310										315					320					
ggt	caa	gga	act	gtg	ttt	gat	gcc	ttc	cgc	gat	cag	gtg	gcc	atc	cag	1124				
Gly	Gln	Gly	Thr	Val	Phe	Asp	Ala	Phe	Pro	Asp	Gln	Val	Ala	Ile	Gln					
			325				330						335							
ctg	aat	gat	act	cac	cct	cgc	atc	gcg	atc	cct	gag	ctg	atg	agg	att	1172				
Leu	Asn	Asp	Thr	His	Pro	Arg	Ile	Ala	Ile	Pro	Glu	Leu	Met	Arg	Ile					
			340				345						350							
ttt	gtg	gat	att	gaa	aaa	ctg	ccc	tgg	tcc	aag	gca	tgg	gag	ctc	aac	1220				
Phe	Val	Asp	Ile	Glu	Lys	Leu	Pro	Trp	Ser	Lys	Ala	Trp	Glu	Leu	Asn					
			355				360						365							
cag	aag	acc	ttc	gcc	tac	acc	aac	cac	aca	gtg	ctc	cgc	gaa	gcc	ctg	1268				
Gln	Lys	Thr	Phe	Ala	Tyr	Thr	Asn	His	Thr	Val	Leu	Pro	Glu	Ala	Leu					
			370				375						380			385				
gag	cgc	tgg	ccc	gtg	gac	ctg	gtg	gag	aag	ctg	ctc	cct	cga	cat	ttg	1316				
Glu	Arg	Trp	Pro	Val	Asp	Leu	Val	Glu	Lys	Leu	Leu	Pro	Arg	His	Leu					
			390							395						400				
gaa	atc	att	tat	gag	ata	aat	cag	aag	cat	tta	gat	aga	att	gtg	gcc	1364				
Glu	Ile	Ile	Tyr	Glu	Ile	Asn	Gln	Lys	His	Leu	Asp	Arg	Ile	Val	Ala					
			405							410						415				
ttg	ttt	cct	aaa	gat	gtg	gac	cct	ctg	aga	agg	atg	tct	ctg	ata	gaa	1412				
Leu	Phe	Pro	Lys	Asp	Val	Asp	Pro	Leu	Arg	Arg	Met	Ser	Leu	Ile	Glu					
			420							425						430				
gag	gaa	gga	agc	aaa	agg	atc	aac	atg	gcc	cat	ctc	tgc	att	gtc	ggt	1460				
Glu	Glu	Gly	Ser	Lys	Arg	Ile	Asn	Met	Ala	His	Leu	Cys	Ile	Val	Gly					
			435				440						445							
tcc	cat	gct	gtg	aat	ggc	gtg	gct	aaa	atc	cac	tca	gac	atc	gtg	aag	1508				
Ser	His	Ala	Val	Asn	Gly	Val	Ala	Lys	Ile	His	Ser	Asp	Ile	Val	Lys					
			450				455						460			465				
act	aaa	gta	ttc	aag	gac	ttc	agt	gag	cta	gaa	cct	gac	aag	ttt	cag	1556				
Thr	Lys	Val	Phe	Lys	Asp	Phe	Ser	Glu	Leu	Glu	Pro	Asp	Lys	Phe	Gln					
			470							475						480				
aat	aaa	acc	aat	ggg	atc	act	cca	agg	cgc	tgg	ctc	cta	ctc	tgc	aac	1604				
Asn	Lys	Thr	Asn	Gly	Ile	Thr	Pro	Arg	Arg	Trp	Leu	Leu	Leu	Cys	Asn					
			485							490						495				
cca	gga	ctt	gca	gag	ctc	ata	gca	gag	aaa	att	gga	gaa	gac	tat	gtg	1652				
Pro	Gly	Leu	Ala	Glu	Leu	Ile	Ala	Glu	Lys	Ile	Gly	Glu	Asp	Tyr	Val					
			500							505						510				
aaa	gac	ctg	agc	cag	ctg	acg	aag	ctc	cac	agc	ttc	ctg	ggt	gat	gat	1700				
Lys	Asp	Leu	Ser	Gln	Leu	Thr	Lys	Leu	His	Ser	Phe	Leu	Gly	Asp	Asp					
			515				520						525							
gtc	ttc	ctc	cgg	gaa	ctc	gcc	aag	gtg	aag	cag	gag	aat	aag	ctg	aag	1748				
Val	Phe	Leu	Arg	Glu	Leu	Ala	Lys	Val	Lys	Gln	Glu	Asn	Lys	Leu	Lys					
			530				535						540			545				
ttt	tct	cag	ttc	ctg	gag	acg	gag	tac	aaa	gtg	aag	atc	aac	cca	tcc	1796				

Phe Ser Gln Phe Leu Glu Thr Glu Tyr Lys Val Lys Ile Asn Pro Ser	
550 555 560	
tcc atg ttt gat gtc cag gtg aag agg ata cat gag tac aag cga cag	1844
Ser Met Phe Asp Val Gln Val Lys Arg Ile His Glu Tyr Lys Arg Gln	
565 570 575	
ctc ttg aac tgt ctg cat gtg atc acg atg tac aac cgc att aag aaa	1892
Leu Leu Asn Cys Leu His Val Ile Thr Met Tyr Asn Arg Ile Lys Lys	
580 585 590	
gac cct aag aag tta ttc gtg cca agg aca gtt atc att ggt ggt aaa	1940
Asp Pro Lys Lys Leu Phe Val Pro Arg Thr Val Ile Ile Gly Gly Lys	
595 600 605	
gct gcc cca gga tat cac atg gcc aaa atg atc ata aag ctg atc act	1988
Ala Ala Pro Gly Tyr His Met Ala Lys Met Ile Ile Lys Leu Ile Thr	
610 615 620 625	
tca gtg gca gat gtg gtg aac aat gac cct atg gtt gga agc aag ttg	2036
Ser Val Ala Asp Val Val Asn Asn Asp Pro Met Val Gly Ser Lys Leu	
630 635 640	
aaa gtc atc ttc ttg gag aac tac aga gta tct ctt gct gaa aaa gtc	2084
Lys Val Ile Phe Leu Glu Asn Tyr Arg Val Ser Leu Ala Glu Lys Val	
645 650 655	
att cca gcc aca gat ctg tca gag cag att tcc act gca ggc acc gaa	2132
Ile Pro Ala Thr Asp Leu Ser Glu Gln Ile Ser Thr Ala Gly Thr Glu	
660 665 670	
gcc tcg ggg aca ggc aat atg aag ttc atg cta aat ggg gcc cta act	2180
Ala Ser Gly Thr Gly Asn Met Lys Phe Met Leu Asn Gly Ala Leu Thr	
675 680 685	
atc ggg acc atg gat ggg gcc aat gtg gaa atg gca gaa gaa gct ggg	2228
Ile Gly Thr Met Asp Gly Ala Asn Val Glu Met Ala Glu Glu Ala Gly	
690 695 700 705	
gaa gag aac ctg ttc atc ttt ggc atg agc ata gat gat gtg gct gct	2276
Glu Glu Asn Leu Phe Ile Phe Gly Met Ser Ile Asp Asp Val Ala Ala	
710 715 720	
ttg gac aag aaa ggg tac gag gca aaa gaa tac tat gag gca ctt cca	2324
Leu Asp Lys Lys Gly Tyr Glu Ala Lys Glu Tyr Tyr Glu Ala Leu Pro	
725 730 735	
gag ctg aag ctg gtc att gat caa att gac aat ggc ttt ttt tct ccc	2372
Glu Leu Lys Leu Val Ile Asp Gln Ile Asp Asn Gly Phe Phe Ser Pro	
740 745 750	
aag cag cct gac ctc ttc aaa gat atc atc aac atg cta ttt tat cat	2420
Lys Gln Pro Asp Leu Phe Lys Asp Ile Ile Asn Met Leu Phe Tyr His	
755 760 765	
gac agg ttt aaa gtc ttt gca gac tac gaa gcc tat gtc aag tgt caa	2468
Asp Arg Phe Lys Val Phe Ala Asp Tyr Glu Ala Tyr Val Lys Cys Gln	
770 775 780 785	

5

gat aaa gtg agt cag ctg tac atg aat cca aag gcc tgg aac aca atg 2516
 Asp Lys Val Ser Gln Leu Tyr Met Asn Pro Lys Ala Trp Asn Thr Met
 790 795 800

gta ctc aaa aac ata gct gcc tgc ggg aaa ttc tcc agt gac cga aca 2564
 Val Leu Lys Asn Ile Ala Ala Ser Gly Lys Phe Ser Ser Asp Arg Thr
 805 810 815

att aaa gaa tat gcc caa aac atc tgg aac gtg gaa cct tca gat cta 2612
 Ile Lys Glu Tyr Ala Gln Asn Ile Trp Asn Val Glu Pro Ser Asp Leu
 820 825 830

aag att tct cta tcc aat gaa tct aac aaa gtc aat gga aat tga 2657
 Lys Ile Ser Leu Ser Asn Glu Ser Asn Lys Val Asn Gly Asn
 835 840 845

actctacaat gtctctagaa aacatagctt cttactgaac ttgaacattt ttacaacatt 2717

cactgggtttt tgttttggtta gctaataatc tataatagtt gagtatctct gggaatgggg 2777

agggaaatta tatgtaatag agcttaaaaa taaagtgtca atttccaagg a 2828

<210> 2
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Primer

<400> 2
 catgggcccga acattacaga a 21

<210> 3
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Primer

<400> 3
 caagaccacc attgccaagt c 21

<210> 4
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Probe

<400> 4
 ctgtgatgag gccatttacc agcttgg 27

6

<210> 5
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 5
gaagggtgaag gtcggagtc

19

<210> 6
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 6
gaagatgggtg atgggatttc

20

<210> 7
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Probe

<400> 7
caagcttccc gttctcagcc

20

<210> 8
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 8
ccgccccgcc gcgccaggag

20

<210> 9
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 9
cgggctgcgc agagagctgg

20

<210> 10
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 10
cagcggttcg cccatggctg 20

<210> 11
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 11
tctgtcagcg gttcgcccac 20

<210> 12
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 12
tgccacgttc tccacgcca 20

<210> 13
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 13
cttgaccagc gtgaagtga 20

<210> 14
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 14
gcgcgaagta gtagtcgcgg 20

<210> 15
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 15
tccagcgccc caccaggtgg

20

<210> 16
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 16
cccatgtaaa attccagaga

20

<210> 17
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 17
atattgcaga ccgaggttga

20

<210> 18
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 18
ctcttctata tccaatccaa

20

<210> 19
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 19

gcaggcagca agtctcccaa

20

<210> 20

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 20

gccgtatcca taggctgcaa

20

<210> 21

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 21

tacctgccat ccatctcgga

20

<210> 22

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 22

gggcgggact tctcccaagg

20

<210> 23

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 23

tcccgggtgtt ggtgtgttct

20

<210> 24

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

10

<400> 24
ggtgttgaca gtgttattca

20

<210> 25
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 25
tccaacatta aagtctctga

20

<210> 26
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 26
attgtcattg ggatagagga

20

<210> 27
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 27
tggagccaaa cttggaggct

20

<210> 28
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 28
ttctggttga gctcccatgc

20

<210> 29
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 29
acggggccagc gctccagggc 20

<210> 30
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 30
atgcttctga tttatctcat 20

<210> 31
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 31
atcagagaca tccttctcag 20

<210> 32
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 32
cacagcatgg gaaccgacaa 20

<210> 33
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 33
gaagtccttg aatacttttag 20

<210> 34
<211> 20
<212> DNA
<213> Artificial Sequence

<220>

12

<223> Antisense Oligonucleotide

<400> 34

ctgcaagtcc tgggttgcag

20

<210> 35

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 35

atcacccagg aagctgtgga

20

<210> 36

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 36

aggatgggtt gatcttcact

20

<210> 37

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 37

acagttcaag agctgtcgct

20

<210> 38

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 38

cagctttacc accaatgata

20

<210> 39

<211> 20

<212> DNA

<213> Artificial Sequence

13

<220>
<223> Antisense Oligonucleotide

<400> 39
ttccaaccat agggtcattg 20

<210> 40
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 40
atctgtggct ggaatgactt 20

<210> 41
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 41
tcatattgcc tgtccccgag 20

<210> 42
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 42
ccacattggc cccatccatg 20

<210> 43
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 43
atgctcatgc caaagatgaa 20

<210> 44
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 44
agtattcttt tgcctcgtag

20

<210> 45
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 45
ttgtcaattt gatcaatgac

20

<210> 46
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 46
tcatgataaa atagcatggt

20

<210> 47
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Antisense Oligonucleotide

<400> 47
ccccattccc agagatactc

20

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/09982

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12Q 1/68; A01N 43/04; C07H 21/04; A61K 31/07

US CL : 435/6, 325, 375, 91.1; 536/24.5, 23.1, 24.3; 514/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 325, 375, 91.1; 536/24.5, 23.1, 24.3; 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6,043,091 A (MONIA et al.) 19 July 1999 (19.07.1999), see claims 1-20.	1-51
Y	NEWARD et al. The polymorphic locus for glycogen storage disease VI (liver glycogen phosphorylase) maps to chromosome 14. Am. J. Hum. Genet., 1987, Vol. 40, pages 351-364, see entire article.	1-51
Y	BURWINKEL et al. Mutations in the liver glycogen phosphorylase gene (PYGL) underlying glycogenosis type VI. Am. J. Hum. Genet., 1998, Vol. 6, pages 785-791.	1-51
Y	MARTIN et al. Discovery of a human liver glycogen phosphorylase inhibitor that lowers blood glucose in vivo. Proc. Natl. Acad. Sci., 1998, Vol. 95, pages 1176-1181.	1-51

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

30 June 2003 (30.06.2003)

Date of mailing of the international search report

06 AUG 2003

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Washington, D.C. 20543-0145

Facsimile No. (703)305-3230

Authorized officer

Valerie Bell-Harrop
Terra C. Gibbs

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/09982

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claim Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
Please See Continuation Sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.: Claims 1-51, SEQ ID NOs. 9, 20, 16, and 31
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

- The additional search fees were accompanied by the applicant's protest.
No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

PCT/US03/09982

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Groups 1-93, Claims 1-51, drawn to an antisense compound 8 to 50 nucleobases in length targeted to a nucleic acid molecule encoding liver glycogen phosphorylase, wherein said compound specifically hybridizes with said nucleic acid molecule encoding liver glycogen phosphorylase and inhibits the expression of liver glycogen phosphorylase, wherein the antisense compound has a sequence comprising SEQ ID NOs. 9-47, 49-66, 75-83, 85-89, 96-116, 122 and 123 and a method of using said antisense compound.

As outlined above, this international searching authority has found 93 inventions claimed in the International Application covered by the claims indicated: Claims 1-51 which specifically claim sequences listed as SEQ ID Nos. 9-47, 49-66, 75-83, 85-89, 96-116, 122 and 123, which are intended to modulate the function and/or expression of liver glycogen phosphorylase.

This international searching authority considers that the international application does not comply with the requirements of unity of invention (Rules 13.1, 13.2 and 13.3) for the reasons indicated below:

The inventions listed as Groups 1-93 do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

According to the guidelines in Section (f)(i)(a) of Annex B of the PCT Administrative Instructions, the special technical feature as defined by PCT Rule 13.2 shall be considered to be met when all the alternatives of a Markus group are of similar nature. For chemical alternatives, such as the claimed antisense sequences, the Markush group shall be regarded as being of similar nature when

(A) all alternatives have a common property or activity and

(B)(1) a common structure is present, i.e., a significant structure is shared by all of the alternatives or

(B)(2) in cases where the common structure cannot be the unifying criteria, all alternatives belong to an art recognized class of compounds in the art to which the invention pertains.

The instant antisense sequences are considered to be each separate inventions for the following reasons:

The sequences do not meet the criteria of (A), common property or activity or (B)(2), art recognized class of compounds. Although the sequence target and modulate expression of the same gene, each antisense sequence behaves in a different way in the context of the claimed invention. Each sequence targets a different and specific region of liver glycogen phosphorylase and each sequence modifies (either increases or decreases) the expression of the gene to varying degrees (per Applicants' Table I in the specification). Each member of the class cannot be substituted, one for the other, with the expectation that the same intended result would be achieved.

Further, although the sequence target the same gene, the sequences do not meet the criteria of (B)(1), as they do not share, one with another, a common core structure. Accordingly, unity of invention between the antisense sequences is lacking and each antisense sequence claimed is considered to constitute a special technical feature.

Applicants will obtain a search of the first sequence listed in the first invention. For every other sequence applicants wish to have searched, applicants need to elect the sequence and pay an additional fee.

INTERNATIONAL SEARCH REPORT

PCT/US03/09982

Continuation of B. FIELDS SEARCHED Item 3:

Biosis, Medline, Caplus, Embase Cancerlit

search terms: liver glycogen phosphorylase, HLGPa, 1, 4 alpha D glucan, orthophosphate alpha D glucosyltransferase, glycogen phosphophorylase liver, EC 2.4.1.1, antisense and ribozyme